



FIG. 1









FIG.30



PRIOR ART

FIG. 4

5/35





FIG.5A







FIG. 6






FIG. 8




FIG.9






FIG. 10A










FIG. 11B

FIG. 12A

FIG. 12B








FIG.13

FINDALPHAS
$$(E_{L+1,1}, E_{L+1,2})$$
:
for  $l \leftarrow L$  downto 2

for  $m \leftarrow 1$  to  $6 \cdot 3^{L-l}$ 

for  $i \in \{0, 2\}$ 

$$\alpha_{l,3m-i}^* = -\frac{(E_{l+1,m}^i)_y^+ + (E_{l+1,m}^i)_y^-}{(E_{l+1,m}^i)_x^+ + (E_{l+1,m}^i)_x^-}$$

$$E_{l,3m-i} = \mathcal{S}(-\alpha_{l,3m-i}) E_{l,3m-i}^i$$

FIG. 14




FIG. 15





FIG.16







FIG. 18







FIG. 21A



FI6, 21B





FIG. 22



.  $T_{eta,eta',\gamma_r}(\gamma')$  and  $ilde{T}_{eta,eta',\gamma_r}(\gamma')$ 




FIG.23C

FIG. 23D

FIG. 24

SPACE

FREQUENCY

~W2

Ðω,






FIG. 25

SPACE





FREQUENCY



25/35

FIG. 26A









FIG. 27A








FIG. 27B





Sampling on the Central Divergent Beam

FIG. 28







FIG. 29 B Separable Resampling:



Resample onto the points of intersection of the original rays with the vertical fans of the final beam

(2) Resample on individual vertical fans (as in the fanbeam case)








FIG. 31




FIG. 32





FIG. 33

FIG. 34



33/35






FIG.35

3.4/35








FIG.36B

